

Supplemental Table of Contents

Supplemental Table 1: ICD 9/10 codes used for cohort and outcome identification

Supplemental Table 2: Adjusted odds ratio for maternal outcomes by PR-AKI status with and without adjustment for preeclampsia

Supplemental Figure 1: Study flow diagram

Supplemental Table 1: ICD 9/10 codes used for cohort and outcome identification.

	NIS	ICD 9	ICD10
Pregnancy, childbirth, and the puerperium	NEOMAT		
ESRD		585.6	N186
Renal transplant patient		V420	Z940
		99681	T8610, T8611, T8612
		5561	OTS00ZZ, OTS10ZZ
		5569	OTY00Z0, OTY00Z1, OTY00Z2, OTY10Z0, OTY10Z1, OTY10Z2
AKI		5845	N17.0
		5846	N17.1
		5847	N17.2
		5848	N17.8
		5849	N17.9
Miscarriage		634.00 to 634.92	O00-O08, O03, O03.9
Preterm		644.00 to 644.21	O60.00, O60.02, O60.03, O47.00, O47.9, O47.02, O47.03, O47.1, O60.10X0, O60.12X0, O60.13X0, O60.14X0
Preeclampsia		642.4 and 642.5	O14
Eclampsia		642.6	O15
Hemorrhage		666.x	O72.x

Supplemental Table 2: Adjusted odds ratio for maternal outcomes by PR-AKI status with and without adjustment for preeclampsia.

Outcomes	Adjusted OR w/o preeclampsia	Adjusted OR with preeclampsia
Miscarriage	1.64 (CI 1.3 – 2.07)	1.71 (1.35 - 2.17)
Preterm labor	0.88 (CI 0.8 - 0.98)	0.84 (0.76 - 0.93)
Pre / Eclampsia	1.07 (CI 0.98- 1.16)	-
Adverse Discharge	1.12 (CI 0.98 -1.27)	1.1 (0.96 – 1.25)
Maternal Mortality	1.53 (CI 1.25-1.88)	1.57 (1.27 - 1.93)

Supplemental Figure 1: Study flow diagram

