


Supplemental Figure 1: Predictive power of fecal calprotectin for identifying mucosal healing at week 52 in patients (AUC=0,863, p=0,005, cut-off: 120,5 µg/g).



Supplemental Table 1: Correlation of 1-year wPCDAI, fecal calprotectin (FC) and C-reactive protein (CRP), with simple endoscopic score for Crohn's disease (SES-CD) subgroups.

| SES-CD | 0-2 (inactive) | 3-6 (mild activity) | 7-15 (moderate activity) | >15 (severe activity) |
|-----------------|-------------------|------------------------|-----------------------------|--------------------------|
| Patients | 13 | 5 | 18 | 4 |
| wPCDAI | 14±8,5 | 21±14 | 28±24 | 43±24 |
| <i>P value</i> | 0,23 | 0,51 | | 0,27 |
| FC (mcg/g) | 131±88 | 230±153 | 399±190 | 616±371 |
| <i>P value</i> | 0,05 | 0,03 | | 0,07 |
| CRP (mg/L) | 8,5±8,2 | 8,8±9,6 | 11,5±9,4 | 16,3±7,4 |
| <i>P value</i> | 0,46 | 0,24 | | 0,06 |

Supplemental Figure 2: Concordance between wPCDAI and Simple Endoscopic Score for Crohn's Disease (SES-CD) according to endoscopic disease severity.

