

Supplementary Table 1: Characteristics of Crohn's disease patients (N=262) grouped by occurrence of drug-related adverse events (AE) versus no drug AE

	With drug AE n (%)	Without drug AE n (%)	Total n (%)	P-value
N (%) patients	77 (29.4)	185 (70.6)	262 (100.0)	
Gender				0.74
Male	42 (54.5%)	105 (56.8%)	147 (56.1)	
Female	35 (45.5%)	80 (43.2%)	115 (43.9)	
Age at diagnosis (years) (median, IQR, range)	12, 5, 1-16	12, 4, 1-17		0.07
Disease duration (years) (median, IQR, range)	4, 3, 0-15	3, 3, 0-13		<0.001
Disease location at diagnosis				0.33
ileal	7 (9.6)	20 (11.5)	27 (10.9)	
colonic	9 (12.3)	29 (16.7)	38 (15.4)	
ileocolonic	53 (72.6)	122 (70.1)	175 (70.9)	
upper GI	4 (5.5)	3 (1.7)	7 (2.8)	
Missing	4	11	15	
IBD family history				0.11
No	70 (90.92)	154 (83.2)	224 (85.5)	
Yes	7 (9.1)	31 (16.8)	38 (14.5)	
EIM				0.12
No	43 (55.8)	122 (66.0)	165 (63.0)	
Yes	34 (44.2)	63 (34.0)	97 (37.0)	
IBD-related surgery				0.29
No	57 (74.0)	148 (80.0)	205 (78.2)	
Yes	20 (26.0)	37 (20.0)	57 (21.8)	

AE, adverse event; IBD, inflammatory bowel disease; CD, Crohn's disease; EIM, extraintestinal manifestation