

Supplementary Table 2: Characteristics of Ulcerative Colitis (N=206) and inflammatory bowel disease unclassified (N=41) patients grouped by occurrence of drug-related adverse events (AE) versus no drug AE

	With drug AE n (%)	Without drug AE n (%)	Total n (%)	P- value
N (%) patients	55 (22.3)	192 (77.7)	247 (100)	
Gender				0.88
Male	27 (49.1)	92 (47.9)	119 (48.2)	
Female	28 (50.90)	100 (52.1)	128 (51.8)	
Age at diagnosis (years) (median, IQR, range)	11, 6, 3-16	12, 6, 1-17		0.74
Disease duration (years) (median, IQR, range)	4, 5, 1-12	4, 3, 0-14		0.16
Disease location at diagnosis				0.44
Pancolitis	36 (75.0)	125 (69.4)	161 (70.6)	
Left-sided colitis	10 (20.8)	37 (20.6)	47 (20.6)	
Proctitis	2 (4.2)	18 (10.0)	20 (8.8)	
Missing	7	12	19	
IBD family history				0.23
No	46 (83.6)	172 (89.6)	218 (88.3)	
Yes	9 (16.4)	20 (10.4)	29 (11.7)	
EIM				0.94
No	43 (78.23)	151 (78.6)	194 (78.5)	
Yes	12 (21.8)	41 (21.4)	53 (21.5)	
IBD-related surgery				0.06
No	49 (89.1)	184 (95.8)	233 (94.3)	
Yes	6 (10.1)	8 (4.2)	14 (5.7)	

AE, adverse event; IBD, inflammatory bowel disease; UC, ulcerative colitis; IBD-U, inflammatory bowel disease unclassified; EIM, extraintestinal manifestation