## **Supplemental Content 11**

 Table. Methods of Reporting Symptoms in Studies Included in the Review (Rated Medium Potential For Bias, With Inclusive Case Definitions and Fixed Time Points)

Reference	Symptoms Reported
Broglio 2007 <sup>1</sup>	Reported as composite.
Broglio 2009 <sup>2</sup>	No comparator.
Cavanaugh	None reported.
2005 <sup>3</sup>	
Collins 1999 <sup>4</sup>	Reported as composite.
Collins 2006 <sup>5</sup>	Reported as composite.
Covassin 2007 <sup>6</sup>	Reported as composite, except for symptoms that associated with gender
	differences.
	Symptoms used in analysis for Key Question 2.
De Monte 2006 <sup>7</sup>	None reported.
Field 2003 <sup>8</sup>	Reported as composite.
Geyer 2009 <sup>9</sup>	No comparator.
Hinton-Bayre	None reported.
1999 <sup>10</sup>	
Iverson 2004 <sup>11</sup>	Reported as composite.
Kontos 2010 <sup>12</sup>	Reported as composite.
Lovell 2003 <sup>13</sup>	Reported as composite.
Lovell 2007 <sup>14</sup>	Reported as composite.
Maddocks	Headache, dizziness, blurred vision, nausea, double vision, noise sensitivity, light
1995 <sup>15</sup>	sensitivity.
McCrea 2002 <sup>16</sup>	None reported.
McCrea 2003 <sup>17</sup>	Reported as composite.
McCrea 2005 <sup>18</sup>	Reported as composite.
Ono 2007 <sup>19</sup>	No comparator.
Papa 2012a <sup>20</sup>	None reported.
Papa 2012b <sup>21</sup>	None reported.
Saadat 2009 <sup>22</sup>	No comparator.
Sim 2008 <sup>23</sup>	None reported.
Smits 2007 <sup>24</sup>	No comparator.
Turedi 2008 <sup>25</sup>	No comparator.
Van Kampen	Reported as composite.
2006 <sup>26</sup>	

## REFERENCES

- 1. Broglio SP, Macciocchi SN, Ferrara MS. Sensitivity of the concussion assessment battery. *Neurosurgery*. Jun 2007;60(6):1050-1057; discussion 1057-1058.
- 2. Broglio SP, Sosnoff JJ, Ferrara MS. The relationship of athlete-reported concussion symptoms and objective measures of neurocognitive function and postural control. *Clinical Journal of Sport Medicine*. Vol 192009:377-382.
- **3.** Cavanaugh J, Guskiewicz K, Giuliani C, Marshall S, Mercer V, Stergiou N. Detecting altered postural control after cerebral concussion in athletes with normal postural stability. *BJSM online*. Nov 2005;39(11):805-811.
- **4.** Collins MW, Grindel SH, Lovell MR, et al. Relationship between concussion and neuropsychological performance in college football players. *Jama*. 1999;282(10):964-970.
- 5. Collins M, Lovell MR, Iverson GL, Ide T, Maroon J. Examining concussion rates and return to play in high school football players wearing newer helmet technology: a three-year prospective cohort study. *Neurosurgery*. 2006;58(2):275-286; discussion 275-286.
- 6. Covassin T, Schatz P, Swanik CB. Sex differences in neuropsychological function and post-concussion symptoms of concussed collegiate athletes. *Neurosurgery*. 2007;61(2):345-350; discussion 350-341.
- 7. De Monte VE, Geffen GM, Massavelli BM. The effects of post-traumatic amnesia on information processing following mild traumatic brain injury. *Brain Inj.* 2006;20(13-14):1345-1354.
- 8. Field M, Collins MW, Lovell MR, Maroon J. Does age play a role in recovery from sports-related concussion? A comparison of high school and collegiate athletes. *J Pediatr*. 2003;142(5):546-553.
- **9.** Geyer C, Ulrich A, Grafe G, Stach B, Till H. Diagnostic value of S100B and neuronspecific enolase in mild pediatric traumatic brain injury. *Journal of Neurosurgery*. *Pediatrics*. 2009;4:339-344.
- **10.** Hinton-Bayre AD, Geffen GM, Geffen LB, McFarland KA, Friis P. Concussion in contact sports: reliable change indices of impairment and recovery. *J Clin Exp Neuropsychol.* 1999;21(1):70-86.
- **11.** Iverson GL, Gaetz M, Lovell MR, Collins MW. Cumulative effects of concussion in amateur athletes. *Brain Inj.* 2004;18(5):433-443.
- **12.** Kontos AP, Elbin RJ 3rd, Covassin T, Larson E. Exploring differences in computerized neurocognitive concussion testing between African American and White athletes. *Archives of Clinical Neuropsychology*. 2010;25:734-744.
- **13.** Lovell MR, Collins MW, Iverson GL, et al. Recovery from mild concussion in high school athletes. *Journal of Neurosurgery*. 2003;98:296-301.
- **14.** Lovell MR, Pardini JE, Welling J, et al. Functional brain abnormalities are related to clinical recovery and time to return-to-play in athletes. *Neurosurgery*. 2007;61(2):352-359; discussion 359-360.
- **15.** Maddocks DL, Dicker GD, Saling MM. The assessment of orientation following concussion in athletes. *Clin J Sport Med.* 1995;5(1):32-35.
- **16.** McCrea M, Kelly JP, Randolph C, Cisler R, Berger L. Immediate neurocognitive effects of concussion. *Neurosurgery*. 2002;50(5):1032-1040; discussion 1040-1032.

- **17.** McCrea M, Guskiesicz KM, Marshall SW, et al. Acute effects and recovery time following concussion in collegiate football players: the NCAA Concussion Study. *Jama*. 2003;290(19):2556-2563.
- **18.** McCrea M, Barr WB, Guskiewicz K, et al. Standard regression-based methods for measuring recovery after sport-related concussion. *J Int Neuropsychol Soc.* 2005;11(1):58-69.
- **19.** Ono K, Wada K, Takahara T, Shirotani T. Indications for computed tomography in patients with mild head injury. *Neurol Med Chir (Tokyo)*. 2007;47(7):291-297; discussion 297-298.
- **20.** Papa L, Lewis LM, Silvestri S, et al. Serum levels of ubiquitin C-terminal hydrolase distinguish mild traumatic brain injury from trauma controls and are elevated in mild and moderate traumatic brain injury patients with intracranial lesions and neurosurgical intervention. *J Trauma Acute Care Surg.* 2012;72(5):1335-1344.
- **21.** Papa L, Lewis LM, Falk JL, et al. Elevated levels of serum glial fibrillary acidic protein breakdown products in mild and moderate traumatic brain injury are associated with intracranial lesions and neurosurgical intervention. *Ann Emerg Med.* 2012;59(6):471-483.
- **22.** Saadat S, Ghodsi SM, Naieni KH, et al. Prediction of intracranial computed tomography findings in patients with minor head injury by using logistic regression. *Journal of Neurosurgery*. 2009;111:688-694.
- **23.** Sim A, Terryberry-Spohr L, Wilson KR. Prolonged recovery of memory functioning after mild traumatic brain injury in adolescent athletes. *Journal of Neurosurgery*. 2008;108:511-516.
- 24. Smits M, Dippel DW, Steyerberg EW, et al. Predicting intracranial traumatic findings on computed tomography in patients with minor head injury: the CHIP prediction rule.[Summary for patients in Ann Intern Med. 2007 Mar 20;146(6):I55; PMID: 17371881]. *Ann Intern Med.* 2007;146(6):397-405.
- **25.** Turedi S, Hasanbasoglu A, Gunduz A, Yandi M. Clinical decision instruments for CT scan in minor head trauma. *J Emerg Med.* 2008;34(3):253-259.
- **26.** Van Kampen DA, Lovell MR, Pardini JE, Collins MW, Fu FH. The "value added" of neurocognitive testing after sports-related concussion. *Am J Sports Med.* 2006;34(10):1630-1635.