
Evaluable cases with urethritis at the STI outpatient clinic with complete known infection status, n=642
Evaluable cases with gonococcal urethritis, n=174 (27%)
Evaluable cases with NGU, n=468 (73%)
Infection with MG, n=105 (16%)
Infection with CT, n=173 (27%)
No infection with NG, CT or MG, n=208 (32%)
Mono NG infection, n=120 (19%)
Dual NG + CT infection, n=14 (2%)
Dual NG + MG infection, n=32 (5%)
Triple NG + CT + MG infection, n=8 (1%)
Mono MG infection, n=87 (14%)
Dual CT + MG infection, n=18 (3%)
Mono CT infection, n=155 (24%)
Clinically not improved, n=8 (7%)
Clinically not improved, n=1 (7%)
Clinically not improved, n=3 (9%)
Clinically not improved, n=3 (38%)
Clinically not improved, n=33 (38%)
Clinically not improved, n=7 (39%)
Clinically not improved, n=24 (15%)
Clinically not improved, n=48 (23%)
MG-WT infection, n=28 (19%)
MG-MRAM infection, n=94 (65%)
Type unknown MG infection, n=23 (16%)
Clinically not improved, n=4 (14%)
Clinically not improved, n=38 (40%)
Clinically not improved, n=4 (17%)