

Supplemental Table S1

Correlation coefficients between the initial and follow-up assessments of the four scales assessing worst and average pain for the Stable sensitivity analysis group (i.e., those reporting “Minimally improved,” “No change,” or “Minimally worse” in the PGIC).

Scale change	Spearman rank correlation coefficients (r_s)	
	Worst pain (n=136)	Average pain (n=136)
VAS (0-10)	.61	.63
VRS-6 (0-5)	.48	.63
NRS-11 (0-10)	.62	.69
FPS-R (0-10, by 2s)	.54	.59

Abbreviations: VAS, Visual Analogue Scale; VRS-6, 6-point Verbal Rating Scale; NRS-11, 11-point Numerical Rating Scale; FPS-R, Face Pain Scale-Revised.

Supplemental Table S2.

Median, interquartile range (IQR), and *P*-values associated with change in pain intensity over time for the four scales assessing worst and average pain for the Stable sensitivity analysis group (i.e., those reporting “Minimally improved,” “No change,” or “Minimally worse” in the PGIC).

	Median (IQR)		P-value
	Baseline	At 4 weeks	
Worst pain intensity (n=136)			
VAS (0-10)	6.55 (4.90, 7.90)	5.95 (4.30, 7.48)	.001
VRS-6 (0-5)	4.00 (3.00, 4.00)	3.00 (3.00, 4.00)	.001
NRS-11 (0-10)	7.00 (5.00, 8.00)	7.00 (5.00, 8.00)	.015
FPS-R (0-10, by 2s)	6.00 (4.00, 8.00)	6.00 (4.00, 8.00)	.384
Average pain intensity (n=136)			
VAS (0-10)	5.05 (3.80, 6.50)	5.10 (3.73, 6.50)	.341
VRS-6 (0-5)	3.00 (3.00, 3.00)	3.00 (3.00, 3.00)	.231
NRS-11 (0-10)	5.00 (4.00, 7.00)	5.00 (4.00, 7.00)	.401
FPS-R (0-10, by 2s)	4.00 (4.00, 6.00)	6.00 (4.00, 6.00)	.114

Abbreviations: VAS, Visual Analogue Scale; VRS-6, 6-point Verbal Rating Scale; NRS-11, 11-point Numerical Rating Scale; FPS-R, Face Pain Scale-Revised.

Supplemental Table S3

Median, interquartile range (IQR), and *P*-values associated with change in pain intensity over time for the four scales assessing worst and average pain for the Improved sensitivity analysis group (i.e., those reporting “Very much improved” or “Much improved” in the PGIC).

	Median (IQR)		P-value
	Baseline	At 4 weeks	
Worst pain intensity (n=88)			
VAS (0-10)	5.20 (3.65, 7.10)	3.40 (1.70, 5.20)	<.001
VRS-6 (0-5)	3.00 (2.00, 4.00)	2.50 (1.00, 3.00)	<.001
NRS-11 (0-10)	5.00 (4.00, 7.00)	4.00 (2.00, 5.00)	<.001
FPS-R (0-10, by 2s)	4.00 (4.00, 6.00)	4.00 (2.00, 4.00)	<.001
Average pain intensity (n=90)			
VAS (0-10)	4.65 (2.35, 6.20)	2.60 (1.30, 4.50)	<.001
VRS-6 (0-5)	3.00 (2.00, 3.00)	2.00 (1.00, 3.00)	<.001
NRS-11 (0-10)	5.00 (3.00, 6.00)	3.00 (2.00, 5.00)	<.001
FPS-R (0-10, by 2s)	4.00 (2.00, 6.00)	2.00 (2.00, 4.00)	<.001

Abbreviations: VAS, Visual Analogue Scale; VRS-6, 6-point Verbal Rating Scale; NRS-11, 11-point Numerical Rating Scale; FPS-R, Face Pain Scale-Revised.

Supplemental Table S4

Median, interquartile range (IQR), and *P*-values associated with change in pain intensity over time for the four scales assessing worst and average pain for the Worsened sensitivity analysis group (i.e., those reporting “Very much worse” or “Much worse” in the PGIC).

	Median (IQR)		P-value
	Baseline	At 4 weeks	
Worst pain intensity (n=26)			
VAS (0-10)	7.15 (4.95, 9.00)	8.00 (7.00, 8.48)	.017
VRS-6 (0-5)	4.00 (3.00, 4.00)	4.00 (3.75, 4.00)	.871
NRS-11 (0-10)	8.00 (5.75, 9.00)	8.00 (7.00, 9.00)	.008
FPS-R (0-10, by 2s)	8.00 (6.00, 8.00)	8.00 (6.00, 8.00)	.169
Average pain intensity (n=24)			
VAS (0-10)	5.60 (3.20, 8.38)	7.55 (6.30, 8.23)	.003
VRS-6 (0-5)	3.00 (3.00, 4.00)	3.00 (3.00, 4.00)	.604
NRS-11 (0-10)	6.00 (5.00, 7.88)	8.00 (6.25, 8.88)	.001
FPS-R (0-10, by 2s)	6.00 (4.50, 7.75)	6.00 (6.00, 8.00)	.054

Abbreviations: VAS, Visual Analogue Scale; VRS-6, 6-point Verbal Rating Scale; NRS-11, 11-point Numerical Rating Scale; FPS-R, Face Pain Scale-Revised.