

TABLE 2 (supplemental digital content 2): Facial sensibility function results for all patients. Q1, Q2, Q3 etc. correspond to the questions of the Facial Sensibility Function Questionnaire (see Figure, Supplemental Digital Content 1). Mean values shown correspond to a likert-type scale between 1-5. Chi-squared analysis reveals no difference in answers between the metopic/unicoronal craniosynostosis (n=15), sagittal craniosynostosis (n=13) and age-matched controls (n=16). P-values shown in bottom row.

	CONTROL (Mean $\pm$ SD) n= 16	SAGITTAL (Mean $\pm$ SD) n= 13	METOPIC/UCS (Mean $\pm$ SD) n= 15	χ^2	p-value
Q1	1.0 $\pm$ 0.0	1.1 $\pm$ 0.3	1.0 $\pm$ 0.0	2.4	0.3
Q2	1.1 $\pm$ 0.5	1.0 $\pm$ 0.0	1.0 $\pm$ 0.0	1.8	0.4
Q3	1.1 $\pm$ 0.3	1.1 $\pm$ 0.3	1.0 $\pm$ 0.0	1.9	0.4
Q4	1.0 $\pm$ 0.0	1.1 $\pm$ 0.6	1.0 $\pm$ 0.0	2.4	0.3
Q5	1.3 $\pm$ 0.6	1.8 $\pm$ 1.3	1.7 $\pm$ 0.9	8.3	0.4
Q6	1.0 $\pm$ 0.0	1.3 $\pm$ 1.1	1.2 $\pm$ 0.4	8.5	0.1
Q7	1.0 $\pm$ 0.0	1.1 $\pm$ 0.3	1.3 $\pm$ 1.0	4.3	0.4
Q8	1.0 $\pm$ 0.0	1.1 $\pm$ 0.3	1.0 $\pm$ 0.0	2.4	0.3
Q9	---	---	---		---
Q10	4.4 $\pm$ 0.9	3.8 $\pm$ 1.3	4.5 $\pm$ 0.5	9.5	0.3
Q11	5.0 $\pm$ 0.0	4.7 $\pm$ 0.6	4.8 $\pm$ 0.4	5.9	0.2
Q12	5.0 $\pm$ 0.0	4.4 $\pm$ 1.2	4.8 $\pm$ 0.6	6.8	0.3