

Table S1: Risk of AMI in whole sample versus sample restricted to participants without missing cholesterol, blood pressure or smoking data

		All participants (original analysis)	No missing cholesterol, blood pressure or smoking data
	HIV status	HR (95% CI)	
No major CVDRF	HIV uninfected	1 (REF)	1 (REF)
	HIV infected	2.0 (1.0-3.9)*	2.0 (1.0-3.8)**
1 major CVDRF	HIV uninfected	2.9 (1.7-4.9)	2.4 (1.4-4.1)
	HIV infected	3.9 (2.3-6.7)	3.5 (2-6.1)
2 major CVDRF	HIV uninfected	4.4 (2.6-7.5)	3.9 (2.3-6.8)
	HIV infected	6.0 (3.4-10.3)	5.8 (3.3-10.2)
3+ major CVDRF	HIV uninfected	5.0 (2.9-8.7)	5.3 (3-9.4)
	HIV infected	7.0 (3.8-12.9)	7.6 (4.1-14.1)
Missing CVDRF	HIV uninfected	2.1 (1.1-3.9)	NA
	HIV infected	3.6 (1.9-7.0)	NA

Figure S1: Age/race-ethnicity adjusted rates of acute myocardial infarction (AMI) by cardiovascular disease risk factor profile (CVDRF) stratified by HIV status with past smokers excluded (see attached)

