

Appendix 1: Questions in the ERACODA Questionnaire regarding functional and mental status three months after COVID-19 diagnosis

Outcome with respect to functional* (Q5.16) and mental** (Q5.17) status after three months:

*Functional status is a patient's ability to perform normal, daily activities such as walking, bathing, dressing, getting out of bed, and using the toilet.

** Mental status is a patient's intellectual capacity, emotional state, and general mental health (including mood, behavior, orientation, judgment, memory, problem-solving ability, and contact with reality).

Q5.16 Did your patient reach his/her pre-COVID-19 functional status three months after initial presentation?

Options: Yes / No / I don't know and could not obtain this information from my colleague(s)
(Please note that the three months time point is important)

Q5.16.1 When do you think your patient will return to his/her pre-COVID-19 physical functional status?

Options: 0 - 3 months / 3 - 6 months / 6-12 months / >1 year / Never
(Additional time beyond the three months time point that this question refers to)

Q5.16.2 What would be the limiting factor (more than one answer may apply)? Options:

- Impairment due to thrombo-embolic events (stroke, pulmonary embolism etc.)
- Impaired lung function (other than caused by pulmonary embolism)
- Decline in cognitive function (other than due to thrombo-embolic event)
- Reduced mobility (other than due to thrombo-embolic event)
- Reduced muscle strength
- Disturbed mental status (anxiety, depression, post-traumatic stress disorder)
- Tiredness
- Other factor(s)

Q5.16.3 Please specify other factor(s) concerning physical functional status:

Q5.17 Did your patient reach his/her pre-COVID-19 mental status three months after initial presentation?

Options: Yes / No / I don't know and could not obtain this information from my colleague(s)
(Please note that the three months time point is important)

Q5.17.1 When do you think your patient will return to his/her pre-COVID-19 mental status?

Options: 0 - 3 months / 3 - 6 months / 6-12 months / >1 year / Never
(Additional time beyond the three months time point that this question refers to)

Q5.17.2. What would be the limiting factor (more than one answer may apply)? Options:

- Delirium
- Depression

- Anxiety
- Bereavement/grief
- Memory loss
- Sleep disturbances
- Post-traumatic stress disorder
- Other factor(s)

5.17.3 Please specify other factor(s) concerning mental status: