

Table 1S. bTCMR cases with their corresponding MC/mm², Area % SR and interstitial inflammation score (i).

Sample number (n=48)	MC/mm ²	Area % SR	Individual Banff Lesion Score (i)
1	76,55	21,43	1
2	46,63	26,82	0
3	34,42	15,43	1
4	31,82	11,34	0
5	28,35	19,32	0
6	24,97	32,36	1
7	23,26	26,21	0
8	18,73	36,43	1
9	17,22	21,42	1
10	16,61	27,49	1
11	12,85	25,20	0
12	12,25	13,01	1
13	11,58	20,17	0
14	11,30	16,59	0
15	11,22	24,03	0
16	10,29	12,15	0
17	9,61	15,90	1
18	9,10	21,22	0
19	8,90	15,15	1
20	8,03	25,27	1
21	7,88	14,49	1
22	7,55	12,38	0
23	6,44	13,90	0
24	5,94	19,98	0
25	5,85	19,84	1
26	5,45	5,76	1
27	4,87	22,99	1
28	4,79	12,20	1
29	4,27	16,35	0
30	4,24	18,22	1
31	4,17	22,30	0
32	3,82	13,28	0
33	3,61	22,73	1
34	3,55	12,47	1
35	3,06	16,59	0
36	3,06	11,24	0
37	2,87	13,61	0
38	2,33	23,35	1
39	1,78	15,01	0
40	1,63	7,34	0
41	1,45	14,26	0
42	1,44	16,75	0

43	1,15	25,78	0
44	1,11	10,49	1
45	0,74	15,72	0
46	0,57	11,72	1
47	0,48	11,02	1
48	0,00	7,64	0

MC/mm² = mast cells found per mm² in the cortical area of the biopsy.

Area % SR = area percentage Sirius Red staining in cortical area of the biopsy.

Individual Banff Lesion Score (i) = interstitial inflammation (i0 = no inflammation or <10% inflammation of unscarred cortical parenchyma; i1 = inflammation in 10 to 25% of unscarred cortical parenchyma).